





令和6年度水稻調査結果

カルパー コシヒカリ 幼穂形成期

1	年次		苗立数	苗立期	1ヵ月後	6月中旬	7月初め	7月中旬	幼穂形成期	出穂10日前	穂揃期	成熟期
調査日	R 6	事務局										
		全組織		5/24	6/2	6/18	7/1		7/7			
	R 5	慣行		5/29	6/7	6/19	7/3		7/10	7/24	8/10	8/30
	平年値	慣行		5/25	6/7	6/19	7/3		7/12	7/26	8/5	8/30
2	組織名		苗立数	苗立期	1ヵ月後	6月中旬	7月初め	7月中旬	幼穂形成期	出穂10日前	穂揃期	成熟期
草丈（cm）	(農)是安営農組合			21.8	27.0				77.9			
	R 6	事務局										
		全組織		21.8	27.0				77.9			
	R 5	慣行		25.1	29.8				74.7		105.9	
	平年値	慣行		20.7	27.9				74.9		111.2	
3	組織名		苗立数	苗立期	1ヵ月後	6月中旬	7月初め	7月中旬	幼穂形成期	出穂10日前	穂揃期	成熟期
茎数（m当）	(農)是安営農組合		23.0	27.0	77.0							
	R 6	事務局										
		全組織	23.0	27.0	77.0							
	R 5	慣行	25.4	48.0	104.0				153.0		115.0	115.0
	平年値	慣行	21.7	29.3	96.0				154.8		112.0	112.0
	目標	なんと	18~24		84~88	105~109	165~180		141.0			105~109
4	組織名		苗立数	苗立期	1ヵ月後	6月中旬	7月初め	7月中旬	幼穂形成期	出穂10日前	穂揃期	成熟期
茎数（m ² 当）	(農)是安営農組合		77	90	257							
	R 6	事務局										
		全組織	77	90	257							
	R 5	慣行	85	160	347				510		383	383
	平年値	慣行	72	98	320				516		373	373
	目標	なんと	60~80		280~290	350~360	550~600		470			350~360
5	組織名		苗立数	苗立期	1ヵ月後	6月中旬	7月初め	7月中旬	幼穂形成期	出穂10日前	穂揃期	成熟期
群落葉色	(農)是安営農組合				4.2	4.4	4.1		4.0			
	R 6	事務局										
		全組織			4.2	4.4	4.1		4.0			
	R 5	慣行					4.0		4.0	3.7	4.4	
	平年値	慣行			4.3	4.3			4.0	4.0	4.3	
	目標	なんと							3.8	3.8	4.2~4.5	



今後の 管理	(幼穂形成期) 幼穂形成期は平年より6日早い7/8頃となっています。 (穂肥) 分施や特別栽培米については、すぐに幼穂長と生育長を確認し、穂肥の施用時期、施用量を判断しましょう。 (水管理) 出穂までは飽水管理を徹底し、稲体の活力維持に努めましょう。 ※生育が早くなっているため、「出穂10日前調査」を7/21(日)に変更します。											調査 次回 日	出穂10日前 7月21日(日)																														
6	組織名		苗立数	苗立期	1ヵ月後	6月中旬	7月初め	7月中旬	幼穂形成期	出穂10日前	穂揃期	成熟期	<table border="1"><caption>Graph Data (Approximate)</caption><thead><tr><th>Date</th><th>R6事務局</th><th>R6全組織</th><th>R5</th><th>平年値</th></tr></thead><tbody><tr><td>5/21</td><td>4.0</td><td>4.0</td><td>4.0</td><td>4.0</td></tr><tr><td>6/10</td><td>7.0</td><td>7.0</td><td>7.0</td><td>7.0</td></tr><tr><td>6/30</td><td>10.0</td><td>10.0</td><td>10.0</td><td>10.0</td></tr><tr><td>7/20</td><td>12.0</td><td>12.0</td><td>12.0</td><td>12.0</td></tr><tr><td>8/9</td><td>13.5</td><td>12.8</td><td>12.8</td><td>13.4</td></tr></tbody></table>	Date	R6事務局	R6全組織	R5	平年値	5/21	4.0	4.0	4.0	4.0	6/10	7.0	7.0	7.0	7.0	6/30	10.0	10.0	10.0	10.0	7/20	12.0	12.0	12.0	12.0	8/9	13.5	12.8	12.8	13.4
Date	R6事務局	R6全組織	R5	平年値																																							
5/21	4.0	4.0	4.0	4.0																																							
6/10	7.0	7.0	7.0	7.0																																							
6/30	10.0	10.0	10.0	10.0																																							
7/20	12.0	12.0	12.0	12.0																																							
8/9	13.5	12.8	12.8	13.4																																							
葉 齢	(農)是安営農組合			4.3	7.0	9.8	11.3		12.0																																		
R 6	事務局																																										
	全組織			4.3	7.0	9.8	11.3		12.0																																		
R 5	慣行			4.8	6.7	8.8	10.3		11.0		12.8																																
平年値	慣行			3.6	6.6	8.7	10.3		11.4		13.4																																
7	組織名		田干し	深水管理	浅水管理	中干し	間断かん水	飽水管理	湛水管理	間断かん水	収穫前落水	8	組織名		中干し前	中干し完了	間断かん水時	幼穂形成期																									
水 管理 (看板)	(農)是安営農組合												(農)是安営農組合			1	1																										
	R 6	事務局											R 6	事務局																													
		全組織												全組織		1	1																										
	R 5	慣行											R 5	慣行																													
平年値	慣行											平年値	慣行																														
目安												目安	移植	15	10	5	1																										

作成中

5/5

5/12

9	組織名		コーティング	調査主体	耕種概要	1ヵ月後	幼穂形成期	出穂10日前	穂揃期	抜き株	坪刈り
調査状況等	(農)是安営農組合		カルパー	生産者	○	○	○				

中干し前

中干し完了

間断かん水時

幼穂形成期

足跡深さ (cm)

0

5

10

15

20

0

5

10

15

20

10	組織名		播種日	幼穂形成期	～	出穂期	～	成熟期
生育時期	(農)是安営農組合		5/1	7/8				
	R 6	事務局						
		全組織	5/1	7/8				
R 5	慣行	5/1	7/11	18	7/29	33	8/31	
平年値	慣行	5/1	7/14	21	8/3	39	9/12	

※ () は予想

11	組織名		播種量	播種深度	m苗立本数	m ² 苗立本数	苗立率
播種状況	(農)是安営農組合		kg/10a	mm	本/m	本/m ²	%
			2.5	5.3	22.3	74.3	81.8
	R 6	事務局					
		全組織	2.5	5.3	22.3	74.3	81.8
R 5	慣行	2.9	15.0	25.4	84.7	81.7	
平年値	慣行	3.0	7.2	21.7	72.4	66.7	
目標	なんと	2.5~3.0	10.0	18~24	60~80	50~80	