





令和6年度水稻調査結果

カルパー てんこもり 成熟期

調査日	1	年次		苗立数	苗立期	1ヵ月後	6月中旬	7月初め	7月中旬	幼穂形成期	出穂10日前	穂揃期	成熟期
	R 6	慣行		5/27	6/3	6/18	7/1		7/16	7/29	8/12		
		全組織		5/27	6/3	6/18	7/1		7/16	7/29	8/12		
	R 5	慣行		5/26	6/7	6/16	7/3		7/18	7/28	8/17	8/30	
		平年値		5/26	6/6	6/18	7/6		7/17	7/30	8/11	8/30	
草丈 (cm)	2	組織名		苗立数	苗立期	1ヵ月後	6月中旬	7月初め	7月中旬	幼穂形成期	出穂10日前	穂揃期	成熟期
	(農)千福営農			13.2	16.0				74.2		105.6		
茎数 (m当)	3	組織名		苗立数	苗立期	1ヵ月後	6月中旬	7月初め	7月中旬	幼穂形成期	出穂10日前	穂揃期	成熟期
	(農)千福営農			17.0	17.0	17.0				159.0		129.0	129.0
茎数 (m当)	4	組織名		苗立数	苗立期	1ヵ月後	6月中旬	7月初め	7月中旬	幼穂形成期	出穂10日前	穂揃期	成熟期
	(農)千福営農			57	57	57				530		430	430
群落葉色	5	組織名		苗立数	苗立期	1ヵ月後	6月中旬	7月初め	7月中旬	幼穂形成期	出穂10日前	穂揃期	成熟期
	(農)千福営農						4.3	4.4			4.2		



黒字を慣行として平均

今後の 管理													調査 次回 日	

葉 齢	6	組織名	苗立数	苗立期	1ヵ月後	6月中旬	7月初め	7月中旬	幼穂形成期	出穂10日前	穂揃期	成熟期	<table><thead><tr><th>日</th><th>R 6 慣行</th><th>R 6 全組織</th><th>R 5 慣行</th><th>平年値</th></tr></thead><tbody><tr><td>5/1</td><td>4.0</td><td>4.0</td><td>3.5</td><td>3.5</td></tr><tr><td>5/21</td><td>6.0</td><td>6.0</td><td>5.0</td><td>5.0</td></tr><tr><td>6/10</td><td>9.0</td><td>9.0</td><td>8.0</td><td>8.0</td></tr><tr><td>6/30</td><td>11.0</td><td>11.0</td><td>10.0</td><td>10.0</td></tr><tr><td>7/20</td><td>13.0</td><td>13.0</td><td>12.0</td><td>12.0</td></tr><tr><td>8/9</td><td>14.0</td><td>15.5</td><td>14.5</td><td>13.5</td></tr></tbody></table>	日	R 6 慣行	R 6 全組織	R 5 慣行	平年値	5/1	4.0	4.0	3.5	3.5	5/21	6.0	6.0	5.0	5.0	6/10	9.0	9.0	8.0	8.0	6/30	11.0	11.0	10.0	10.0	7/20	13.0	13.0	12.0	12.0	8/9	14.0	15.5	14.5	13.5
	日	R 6 慣行	R 6 全組織	R 5 慣行	平年値																																											
	5/1	4.0	4.0	3.5	3.5																																											
	5/21	6.0	6.0	5.0	5.0																																											
	6/10	9.0	9.0	8.0	8.0																																											
	6/30	11.0	11.0	10.0	10.0																																											
	7/20	13.0	13.0	12.0	12.0																																											
	8/9	14.0	15.5	14.5	13.5																																											
	(農)千福営農		4.1		9.9	12.4																																										
R 6 慣行		4.1		9.9	12.4																																											
全組織		4.1		9.9	12.4																																											
R 5 慣行		3.2	6.7	8.7	11.7		13.6		15.3																																							
平年値		3.6	6.9	9.1	11.1		12.7		14.6																																							

調 査 状 況 等	7	組織名	ローティング	調査主体	慣行	1ヵ月後	幼穂形成期	出穂10日前	穂揃期	成熟期	抜き株	坪刈り	採取日	指摘事項
	(農)千福営農	カルパー	生産者	慣行	○	○	○	○				○	9/20	抜き株サンプル無い

天候

上旬	中旬	下旬	上旬	中旬	下旬	上旬	中旬	下旬	上旬	中旬	下旬	上旬	中旬	下旬
5月			6月			7月			8月			9月		

水 管 理 看 板	8	組織名	田干し	深水管理	浅水管理	中干し	間断かん水	飽水管理	湛水管理	間断かん水	収穫前落水	足 跡 深 さ	9	組織名	中干し前	中干し完了	間断かん水時	幼穂形成期
	(農)千福営農	4/30		5/17	6/8	6/26							(農)千福営農		10	3		

生 育 時 期	10	組織名	播種日	幼穂形成期	～	出穂期	～	成熟期	播 種 状 況	11	組織名	播種量	播種深度	m苗立本数	kg苗立本数	苗立率
	(農)千福営農	4/30	7/19	20	8/8	40	9/17	(農)千福営農		3.0	4.9	17.3	57.7	52.9		


R 6 慣行	4/30	7/19	20	8/8	40	9/17	R 6 慣行	3.0	4.9	17.3	57.7	52.9	
全組織	4/30	7/19	20	8/8	40	9/17	全組織	3.0	4.9	17.3	57.7	52.9	
R 5 慣行	5/3	7/18	22	8/9	39	9/17	R 5 慣行	7.5	27.2	90.7	83.1		
平年値	4/30	7/18	21	8/8	44	9/21	平年値	5.9	26.3	87.6	75.2		
							目標	なんと	2.7~3.0	10.0	18~24	60~80	50~80

※ () は予想


※ ○ は予想

カルパー	てんこもり	成熟期
------	-------	-----


12	組織名	全重	ワラ重	初重	初ワラ比	粗立米重	精立米重	屑米重	毎トラ歩合
		kg/10a	kg/10a	kg/10a		kg/10a	kg/10a	kg/10a	%
	(農)千福宮農	1970	1037	871	0.8	715	680	35	78.1
R 6	慣行	1970	1037	871	0.8	715	680	35	78.1
	全組織	1970	1037	871	0.8	715	680	35	78.1
R 5	慣行	1755	909	787	0.9	643	624	18	79.3
平成値	慣行	1711	886	781	0.9	648	593	55	76.9



13	組織名		㎡当穂数	1穂初数	㎡当初数	登熟歩合	千粒重	理論収量
			本/㎡	粒/穂	百粒/㎡	%	g	kg/10a
	(農) 千富宮農		430				22.0	
収量構成要素	R 6	慣行	430				22.0	
		全組織	430				22.0	
	R 5	慣行	603	68	408	88.8	22.2	805
	平年値	慣行	492	71	347	78.5	22.3	606
	目標	みんし	420	75	315	80.0	22.5	570




粒厚分布（%）	14	組織名	2.2mm以上	2.2°2.1	2.1°2.0	2.0°1.9	1.9°1.85	1.85°1.8	1.8°1.7	1.7mm以下	合計	1.9mm以上	1.85mm以下
		(農)千福富農	1.4	15.0	57.4	21.4	2.7	0.8	0.9	0.6	100	95.1	97.8
	R 6	慣行	1.4	15.0	57.4	21.4	2.7	0.8	0.9	0.6	100	95.1	97.8
		全組織	1.4	15.0	57.4	21.4	2.7	0.8	0.9	0.6	100	95.1	97.8
	R 5	慣行	2.7	28.3	51.5	14.7	1.4	0.3	0.6	0.4	100	97.2	98.6
		平年値	3.0	21.9	42.8	24.9	3.1	1.4	1.5	1.4	100	91.7	95.9



黒字を慣行として平均

節間長・葉身長等	15	組織名		穂長	第1節間	第2節間	第3節間	第4節間	第5節間	第6節間	合計	葉身長1	葉身長2	葉身長3	倒伏割合(%)			倒伏率		
			cm	cm	cm	cm	cm	cm	cm	cm	cm	cm	cm	cm	少	中	多	甚	%	
		(農)千福宮農																		
		R 6	慣行																	
			全組織																	
	R 5	慣行	18.7	36.7	18.5	14.5	9.6	4.0	0.4	83.7	23.7	34.9	42.9							
	並年値	慣行	18.5	36.5	19.6	13.9	9.0	3.7	0.2	82.9	25.8	36.1	42.9							



16	組織名		整粒	乳白粒	青未熟	基部	腹白	他未計	発芽粒	胴割粒	奇形粒	茶米	斑点粒	タンパク	粒長	粒幅
			%	%	%	%	%	%	%	%	%	%	%	%	mm	mm
	(農) 千福営業		82.6	0.1	0.5	0.2	0.1	15.4	0.1	0.5	0.2	0.2	0.0	5.9	5.25	2.9
外観品質・食味等	R 6	慣行	82.6	0.1	0.5	0.2	0.1	15.4	0.1	0.5	0.2	0.2	0.0	5.9	5.25	2.91
		全組織	82.6	0.1	0.5	0.2	0.1	15.4	0.1	0.5	0.2	0.2	0.0	5.9	5.25	2.91
	R 5	慣行	87.4	0.5	0.7	0.3	0.3	10.4	0.0	0.1	0.0	0.1	0.1	6.1		
	平成28年産	慣行	83.6	0.5	2.0	0.4	0.2	11.3	0.0	0.1	0.0	0.0	0.1	6.3	5.14	3.0